arXiv:1211.2993v1 [quant-ph] 13 Nov 2012

A quantum non-Gaussian Fock state from a semiconductor quantum dot

Ana Predojević[1], Miroslav Ježek[2], Tobias Huber[1], Harishankar Jayakumar[1],
Thomas Kauten[1], Glenn S. Solomon[3], Radim Filip[2], and Gregor Weihs[1]
[1] *Institute for Experimental Physics, University of Innsbruck, Technikerstrasse 25, 6020 Innsbruck, Austria*
[2] *Department of Optics, Palacký University, 17. listopadu 12, 77146 Olomouc, Czech Republic and*
[3] *Joint Quantum Institute, National Institute of Standards and Technology and University of Maryland, Gaithersburg, MD 20849, USA*

Quantum dots are single photon emitters [1]. A single photon state is quantum mechanically represented by the Fock state, $|1\rangle$, which is a quantum non-Gaussian state characterized by a Wigner function negative at the origin [2]. Due to the low collection and detection efficiencies, the reconstruction of the Wigner function from measurements of the quantum dot emission would not confirm the non-Gaussian nature of the emitted light since the negativity of the Wigner function is lost for detection efficiencies below 50%. Semiconductor photon sources like quantum dots are commonly characterized as single photon sources through an auto-correlation measurement [3]. In this work instead, we use a criterion [4] based on an estimate of the photon-number distribution. Through our measurements we prove that the state emitted from a resonantly excited quantum dot is unambiguously a non-Gaussian state.

The quantum nature of a single photon is both a fundamental witness of the quantum character of the light generating process, as well as an important operational resource for quantum information science [5]. Single photons produced by matter are an evidence of quantum jumps; only a discrete level structure can produce such highly non-classical states of light. Traditionally, the boundary between classical and quantum states of light is formed by mixtures of coherent states. Coherent states of light are most similar to classical waves and they can be produced, in principle, by a classical continuous driving of the quantum oscillator through a linear interaction [6]. Such generation processes are governed by a quantum interaction Hamiltonian being linear in the annihilation and creation operators. To recognize that the light comes from a source incompatible with only linear interactions, one can perform the photon auto-correlation experiment [3], which uses the indivisibility of a single photon. This method is accepted as a common way to test a light source for non-classicality [7].

A more demanding level of witnessing the quantum character of light is represented by all possible linear and quadratic Hamiltonian operators. The processes described by these operators can produce any mixture of Gaussian states of light including, for example, the squeezed states of light [8–10]. A mixture of Gaussian states of a single mode of light can be described as a density matrix $\rho = \int P(\lambda)|\lambda\rangle\langle\lambda| d\lambda$, where $|\lambda\rangle$ is a pure Gaussian state of light parametrized by the multi-index λ of its complex amplitude α, squeezing, and phase between the amplitude and direction of squeezing [11]. $P(\lambda)$ is a distribution function. For the auto-correlation test, $|\lambda\rangle$ can be reduced to the coherent state and $P(\alpha)$ is then the traditional Glauber-Sudarshan quasi-distribution [6]. Although the auto-correlation is a valid test of the nonclassicality of the emitted state, it is not sufficient to prove its non-Gaussian nature since an arbitrarily small value of the second-order correlation function can also be achieved by Gaussian states [12]. Consequently, it is important to exclude the possibility that a photon source is governed only by quantum processes up to quadratic order and classical non-linear mechanisms that can mix Gaussian states of light.

Recently, a novel unconditional criterion of quantum non-Gaussianity suitable for single photon sources has been proposed [4] and successful proof-of-principle tests have been performed on a parametric down-conversion source with conditional single photon state preparation [13, 14]. This criterion can detect the non-Gaussianity of sources exhibiting very low collection and detection efficiency without any corrections. Therefore, this criterion is very suitable for the detection of a single photon nature of a solid-state quantum emitter. Both single photon and homodyne detectors are capable of meeting this criterion. The criterion remains valid for multimode states; moreover, it is extremely more sensitive to a multimode generation process than the auto-correlation test [4]. In comparison to the auto-correlation test, this criterion gives a more complete description of the statistics of light. It uses a lower bound of the estimated probability p_1 of successful single photon generation for fixed probability p_{2+} of error representing the generation of two or more photons. The theory of the criterion is detailed in [4, 13].

Here, we present measurements performed to test the non-Gaussian nature of the emission of a quantum dot. To drive our quantum dot system we used two types of excitation: above-band and resonant. In above-band excitation the energy of the excitation laser is much larger than the quantum dot emission energy. It produces carriers in the surrounding material that can be randomly trapped in the quantum dot potential. Secondly, we performed resonant excitation [15] by two-photon excitation of the biexciton state. Here, we exploited the biexciton binding energy [15, 16] in order to use an excitation laser not resonant with any single photon transition. With this laser we drove the two-photon resonance via a virtual level placed energy-wise half way between the exciton and biexciton photon energies. The scheme of the energy levels and the photo-luminescence (PL) of the quantum dot under resonant excitation are shown in Fig. 1 and the experimental set-up is shown in Fig. 2. The resonant excitation scheme gives an excellent suppression of multi-photon events, as evident in Fig. 3a.

To estimate the photon statistics, we firstly derive the number of single counts, two-fold, and three-fold coincidences from the measured data, which includes the arrival times of all the recorded photon detection events. Here, single counts are detections of a biexciton photon (trigger event). A two-fold coincidence is the detection of a biexciton photon followed by a detection of an exciton photon in one of the arms of the beamsplitter. A three-fold coincidence stands for a detection event on all three detectors. The coincidence window was varied as detailed below.

As described in [13] we used the number of single counts, two-fold, and three-fold coincidences to estimate the contribution of vacuum p_0, single photon p_1, and multi-photon terms p_{2+} (see supplementary information), which form the emitted (exciton) signal. From these quantities (probabilities) we calculated the non-Gaussianity witness, ΔW, using the method described in [4, 13] and supplementary information. The results are shown in Fig. 3b and Tables I and II. The values of the witness $\Delta W > 0$ indicate that the measured state is non-Gaussian. Due to the very low collection efficiency ($\approx$ 0.27%) the vacuum term p_0 prevails. The statistical uncertainties for p_0, p_1, and p_2+ were determined from the Poissonian statistics of the recorded events. In Fig. 3b, the states of light in the white area are non-Gaussian and produced by a source incompatible with only quadratic non-linear processes. For the resonantly excited quantum dot our results unambiguously prove that the state cannot be expressed as a mixture of Gaussian states, because the witness, ΔW is positive for any coincidence window that is smaller than the repetition rate of the laser pulses. For a coincidence window of 5.12 ns the measured state exceeds the Gaussian boundary by one standard deviation (green circle in Fig. 3b). Extending the coincidence window to 12.29 ns, which includes the beginning of the consecutive pulse, we find the measured state at the boundary of the Gaussian states. Further extension of the coincidence window places the state in the region where we cannot distinguish it from a coherent mixture of Gaussian states (yellow circles). The results obtained in above-band excitation show a tendency of approaching the boundary but cannot exceed it due to limitations of the measurement system, as detailed below.

Due to the low collection and detection efficiencies the measurement time had to be considerably long to reach the presented statistical confidence, which led to significant noise in the detected signal. For example, in continuous wave measurements, given the detector dark count rates, the measurement duration, and the single count detection rates we had a noise floor of 0.41 three-fold coincidences for a coincidence window of 1.54 ns. This



FIG. 1: a) Excitation level scheme. Resonant excitation coherently drives the two-photon transition between the ground $|g\rangle$ and the biexciton $|b\rangle$ state via a virtual level shown in dashed gray line. The system decays as a cascade via the exciton $|x\rangle$ state. Of the two possible decay paths we use only the shown vertical polarization. Above-band excitation excites the carriers in the surrounding material. b) We exploit the biexciton binding energy to resonantly excite the quantum dot. The laser wavelength is two-photon resonant with the biexciton energy, and therefore placed in the middle between the exciton and biexciton line.



FIG. 2: Experimental set-up. Light derived from a tunable laser is spectrally filtered in a pulse-stretcher and directed via a single mode fibre onto the quantum dot sample. The polarization of the excitation light is chosen to be horizontal. The emission is collected from the top using a high NA objective. Emission is firstly spectrally resolved on a diffraction grating and the lines of interest (exciton and biexcion) are separated and coupled into optical fibres. A fibre beamsplitter divides the exciton light onto two detectors for state verification. The biexciton detections are used as a trigger event.

TABLE I: Above-band excitation estimated probabilities p_0, p_1, p_{2+} and the corresponding witness ΔW are shown for several different coincidence window widths w.

w [ns]	p_0	p_1 [$\times 10^{-3}$]	p_{2+} [$\times 10^{-8}$]	ΔW [$\times 10^{-8}$]
1.536	0.997553(6)	2.446(6)	6.92 ± 4.89	-5.9
2.048	0.997140(7)	2.859(7)	38.46 ± 11.59	-37
2.56	0.996885(7)	3.114(7)	80.55 ± 16.78	-79
3.072	0.996660(7)	3.339(7)	119.2 ± 20.4	-117
3.84	0.996319(8)	3.678(8)	231.4 ± 28.5	-228

TABLE II: Resonant pulsed excitation allows us to distinguish our state from a mixture of Gaussian states, which is witnessed by $\Delta W > 0$.

w [ns]	p_0	p_1 [$\times 10^{-3}$]	p_{2+} [$\times 10^{-8}$]	ΔW [$\times 10^{-8}$]
5.12	0.997287(3)	2.712(3)	0.66 ± 0.66	0.67
12.29	0.997284(3)	2.715(3)	1.32 ± 0.93	0.012
12.54	0.997279(3)	2.721(3)	2.64 ± 1.32	-1.30

noise floor does not allow the detection of non-Gaussian states of light in the given measurement. The measurement limit of the above-band excitation measurement, which is induced by the dark counts, is depicted as orange dot-dashed line in the figure Fig. 3b. The analysed results obtained under continuous wave excitation were calculated for coincidence windows longer than the lifetime of the exciton state (0.71 ns) where the system can be re-excited. For shorter coincidence windows we do not observe any three-fold coincidences. However, without detecting any three-fold coincidences we cannot build a statistical confidence.

For resonant excitation we estimate the noise floor to be 0.0024 three-fold coincidences due to two reasons: different detectors with lower dark count rate of about 100 counts per second and the different nature of the excitation. Resonant excitation produces predominantly cascaded emissions therefore the heralding efficiency is, even for much lower exciton detection rates, comparable with above-band excitation.

In conclusion, we have demonstrated the non-Gaussian nature of the emission of a single quantum dot under resonant excitation. With this we have detected a higher order non-classicality than usually detected by auto-correlation measurements. Therefore, we gain an intrinsically higher sensitivity to possible contribution from other emitters [4]. Compared with above-band excitation, driving of the quantum dot resonantly showed exceptionally pure quantum states of light. To our knowledge, this measurement is the first demonstration of the non-Gaussian nature of photons produced by a semiconductor device. With increased detection efficiencies available in different device geometries [18] the criterion we used becomes an important measure of the quality of the produced light.

Acknowledgements

This work was funded by the European Research Council (project EnSeNa) and the Canadian Institute for Advanced Research through its Quantum Information Processing program. A.P. would like to thank the Austrian Science Fund (FWF) for the support provided through Lise Meitner Postdoctoral Fellowship M-1243. G.S.S. acknowledges partial support through the Physics Frontier Center at the Joint Quantum Institute



FIG. 3: a) The auto-correlation of the exciton signal shows excellent suppression of multi-photon events which can be quantitatively expressed in $g^{(2)}(0)$=0.0073(8). The plotted data was acquired without the triggering on biexciton photon and is presented without background subtraction. The decaying peak height observable on both sides of the graph results from the blinking of the quantum dot [17]. b) The multiphoton contribution p_2+ is plotted as a function of the single photon contribution, p_1. Here, $\mathcal{G}$ is the set of all mixtures of Gaussian states, and the lower, white region indicates non-Gaussian states. The circles stand for results obtained in resonant and pulsed excitation while triangles for above-band and continuous wave excitation. The error bars represent standard deviations, the horizontal error bars of p_1 are smaller than the size of the symbols. The solid blue curve represents the boundary given by Eq. 5 of the supplementary information. The orange dot-dashed line marks the limit of the detection system in continuous excitation.

(PFC@JQI). R.F. acknowledges project P205/12/0577 of GAČR.

Methods

The sample contained a low density self-assembled InAs quantum dots embedded in planar a micro-cavity. It was placed in a continuous-flow cryostat and held at a temperature of 5K. The excitation light was derived from a 76 MHz tunable Ti:Sapphire laser. When the selected quantum dot was excited resonantly we used a pulse stretcher with spectral filtering to optimize the spectral width of the pulses. In continuous wave excitation this step was not necessary. The light was directed onto a dot from the side, where we used the lateral waveguiding mode of a planar micro-cavity. The emission was collected from top and sent into a home-made grating spectrometer. The exciton and biexciton photons were separated by the grating and sent into optical fibres. For continuous wave measurements we used a single mode fibre for the biexciton line and, in order to increase the detection efficiency, a multi-mode fibre beamsplitter for the exciton line. Under resonant excitation it was not possible to use multi-mode fibres due to the spectral proximity of the scattered laser. Measurements were performed using three single photon detectors, two detecting the exciton photons and one detecting biexciton photons. Photon detection events were recorded by a multichannel event timer.

We made two series of measurements with different experimental parameters. The measurements of the first type were continual and performed using continuous wave above-band excitation. The overall measurement time was close to eight hours and the average count rates in the heralding (biexciton) channel S_0 and the signal (exciton) channels S_{1A} and S_{1B} were {5.4, 358, and 196} kcounts per second, respectively. The beamsplitter ratio for these measurements was T_{mm}=0.64(1). The estimate of the beamsplitter ratio [13] was taken into account in the data processing. The detector dark count rate was 500 counts per second. The second type of measurement was one continuous measurement where the dot was excited pulsed and resonantly. Here the measured rates were { S_0, S_{1A} and S_{1B} } = {23, 10, and 9} kcounts per second with the respective estimate of the beamsplitter ratio of T_{sm}=0.54(1). The measurement time was four hours.

Supplementary material

Here, R_0 is the total number of counts (”singles”) in the biexciton signal for the given measurement time. The two-fold coincidences R_{1A} and R_{1B} as well as the three-fold coincidences R_2 were analysed from measurements. It is shown in [13] that the probability of no event p_0 can be expressed as

$$p_0 = 1 - \frac{R_{1A} + R_{1B} + R_2}{R_0}. \tag{1}$$

The lower bound estimator of the single photon contribution is given by

$$p_1 = \frac{R_{1A} + R_{1B}}{R_0} - \frac{T^2 + (1-T)^2}{2T(1-T)} \frac{R_2}{R_0}, \tag{2}$$

where T is the splitting ratio of the beamsplitter used in the measurements. The contribution p_{2+} of the multiphoton terms can be estimated as

$$p_{2+} = 1 - p_0 - p_1. \tag{3}$$

The criterion introduced in [13] defines a witness,ΔW that the given state is not a mixture of Gaussian states $\rho \notin \mathcal{G}$, where $\mathcal{G}$ is the set of all mixtures of Gaussian states. The non-Gaussianity witness is calculated [13] as

$$W(a) = ap_0 + p_1 \tag{4}$$

where the probabilities p_0 and p_1 are obtained experimentally. If $W(a) > W_G(a)$ then $\rho \notin \mathcal{G}$, where $W_G(a)$ is a tangent on the boundary between physical states which can be described as a mixture of Gaussian states and those that cannot. This bound was given in [13] as

$$p_0 = \frac{e^{-d^2[1-\tanh(r)]}}{\cosh(r)}, \qquad p_1 = \frac{d^2\, e^{-d^2[1-\tanh(r)]}}{\cosh^3(r)}, \tag{5}$$

where r is the squeezing constant and d is the displacement. The witness, ΔW is expressed as the distance between $W(a)$ and $W_G(a)$.

[1] Michler, P. *et al.* A Quantum Dot Single-Photon Turnstile Device. *Science* **290**, 2282-2285 (2000).

[2] Lvovsky, A. I. *et al.* Quantum State Reconstruction of the Single-Photon Fock State. *Phys. Rev. Lett.* **87**, 050402-050405 (2001).

[3] Grangier, P., Roger, G. and Aspect, A. Experimental evidence for a photon anticorrelation effect on a beam splitter : a new light on single-photon interferences. *Europhys. Lett.* **1**, 173-179 (1986).

[4] Filip, R. and Mišta, L. Jr. Detecting Quantum States with a Positive Wigner Function beyond Mixtures of Gaussian States. *Phys. Rev. Lett.* **106**, 200401-200404 (2011).

[5] Santori, C., Fattal, D. and Yamamoto, Y. *Single-photon Devices and Applications.* (Wiley-VCH, Weinheim, 2010).

[6] Glauber, R. *Quantum Theory of Optical Coherence.* (Wiley-VCH, Weinheim, 2007).

[7] Scheel, S. Single-photon sources: an introduction. *Journal of Modern Optics* **56**, 141-160 (2009).

[8] Ou, Z. Y., Pereira, S. F. and Kimble, H. J. Quantum noise reduction in optical amplification. *Phys. Rev. Lett.* **70**, 32393242 (1993).

[9] Vahlbruch, H. *et al.* Observation of Squeezed Light with 10-dB Quantum-Noise Reduction. *Phys. Rev. Lett.* **100**, 033602-033605 (2008).

[10] Takeno, Y., Yukawa, M., Yonezawa, H. and Furusawa, A. Observation of -9 dB quadrature squeezing with improvement of phase stability in homodyne measurement. *Optics Express* **15**, 4321-4327 (2007).

[11] Braunstein, S.L. and van Loock, P. Quantum information with continuous variables. *Rev. Mod. Phys.* **77**, 513577 (2005).

[12] Stoler, D. Photon Antibunching and Possible Ways to Observe It. *Phys. Rev. Lett.* **33**, 13971400 (1974).

[13] Ježek, M., Straka, I., Mičuda, M., Dušek, M., Fiurášek, J. and Filip, R. Experimental Test of the Quantum Non-Gaussian Character of a Heralded Single-Photon State. *Phys. Rev. Lett.* **107**, 213602-213605 (2011).

[14] Ježek, M. *et al.* Experimental test of the strongly nonclassical character of a noisy squeezed single-photon state. *Phys. Rev. A* **86**, 043813 (2012).

[15] Jayakumar, H. *et al.* Deterministic Photon Pairs via Coherent Optical Control of a Single Quantum Dot. arXiv:1211.2613 (2012).

[16] Flissikowski, T. *et al.* Two-Photon Coherent Control of a Single Quantum Dot. *Phys. Rev. Lett.* **92**, 227401-227404 (2004).

[17] Santori C. *et al.* Submicrosecond correlations in photoluminescence from InAs quantum dots. *Phys. Rev. B.* **69**, 205324-205331(2004).

[18] Dousse, A. *et al.* Ultrabright source of entangled photon pairs. *Nature* **466**, 217-220 (2010).